

2014

( 5th Semester )

PHYSICS

SIXTH PAPER

( Quantum Mechanics—II )

Full Marks : 75

Time : 3 hours

( PART : B—DESCRIPTIVE )

( Marks : 50 )

*The figures in the margin indicate full marks  
for the questions*

1. (a) Obtain de Broglie relation by using Lorentz transformation and the standard wave equation. 7
- (b) What do you mean by quantum numbers? Write down the possible quantum numbers for  $n = 2$ . 3

Or

- (a) Show that de Broglie wavelength for a material particle of rest mass  $m_0$  and

G15—250/134a

( Turn Over )

charge  $q$ , accelerated from rest through a potential difference of  $V$  volts relativistically is given by

$$\lambda = \frac{h}{\sqrt{2m_0qV\left(1 + \frac{qV}{2m_0c^2}\right)}} \quad 5$$

- (b) Show that material particle can only be represented by a group wave, not by single wave. 5
2. (a) Show that two eigenfunctions of a Hermitian operator belonging to two distinct eigenvalues are orthogonal. 5
- (b) Show that if two Hermitian operators commute, their product is also Hermitian. 5

Or

- (a) What do you mean by basis set? Do the following matrices form a basis set for a vector space of  $2 \times 2$  matrices?

$$M_1 = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, \quad M_2 = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \quad M_3 = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}, \quad M_4 = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}$$

Justify your answer. 5

- (b) Let  $|\psi\rangle = 2|u_1\rangle - 3|u_2\rangle + i|u_3\rangle$  and  $|\phi\rangle = 3|u_1\rangle - 2|u_2\rangle + 4|u_3\rangle$  and a constant  $a = 3 + 3i$ . Compute the inner product  $\langle\psi|\phi\rangle$  and  $|\alpha\psi\rangle$ . 5

3. (a) Obtain the time-independent Schrödinger equation. 6

(b) The wave function of a particle is given by  $\psi(x) = Ae^{-a^2x^2}$ ,  $-\infty < x < +\infty$ . Obtain the expression for normalization constant and the probability of finding the particle in the region  $0 < x < \infty$ . 4

Or

(a) Obtain the equation for conservation of probability in quantum mechanics. Write the physical meaning of the equation. 6

(b) What do you mean by expectation values in quantum mechanics? Show that the expectation value of Hamiltonian  $H$  is the total energy of the system. 1+3=4

4. What do you mean by Quantum mechanical tunneling effect? Show that the transmittance of a particle incident at rectangular potential barrier is given by

$$T = \frac{16E(V_0 - E)}{V_0^2} e^{-\frac{2\sqrt{2m(V_0 - E)}}{h}a}$$

where  $V_0$  is the potential barrier,  $a$  is barrier thickness. 10

Or

A free particle of energy  $E$  is incident on a potential step given by  $V=0$ ;  $x < 0$  and  $V=V_0$ ;  $x \geq 0$ . Obtain the expressions for transmittance and reflectance for the case  $E > V_0$ .

5. (a) Write down Pauli spin matrices and show that  $[\sigma^2, \sigma_x] = 0$ . 1+4=5
- (b) Show that—
- (i)  $\sigma_x \sigma_y + \sigma_y \sigma_x = 0$ ;
- (ii)  $[\sigma_x, \sigma_y] = 2i\sigma_z$ . 2+3=5

Or

- (a) Show that commutation relation between  $X$  and  $Y$  components of angular momentum  $L_x$  and  $L_y$  is given by  $[L_x, L_y] = i\hbar L_z$ . 5
- (b) What do you mean by orbital gyromagnetic ratio for an electron? Obtain the expression for it. 1+4=5

\*\*\*

2014

( 5th Semester )

**PHYSICS**

SIXTH PAPER

**( Quantum Mechanics—II )**

( PART : A—OBJECTIVE )

( Marks : 25 )

*The figures in the margin indicate full marks for the questions*

SECTION—I

( Marks : 10 )

Put a Tick (✓) mark against the correct answer in the brackets provided : 1×10=10

1. Of the following particles moving with the same velocity, the one which has the largest wavelength is

(a) an electron ( )

(b) a proton ( )

(c) a neutron ( )

(d) an  $\alpha$ -particle ( )

2. According to Schrödinger, a particle is equivalent to

- (a) a single wave ( )
- (b) a wave packet ( )
- (c) a light wave ( )
- (d) Cannot behave as wave ( )

3. Eigenvalues of Hermitian operators

- (a) are real only ( )
- (b) are imaginary only ( )
- (c) can be real or imaginary ( )
- (d) are always complex ( )

4. Let  $u = (1, 2, 4)$  and  $v = (2, -3, 5)$  be any two vectors in  $R^3$  space. Their inner product is equal to

- (a) 10 ( )
- (b) 7 ( )
- (c) -9 ( )
- (d) 16 ( )

5. Let  $\psi$  be a wave function, the quantity  $\psi^*\psi$  represents

- (a) probability density ( )  
 (b) charge density ( )  
 (c) energy density ( )  
 (d) wave intensity ( )

6. Conservation of probability in quantum mechanics is represented by the equation

(a)  $\frac{\partial \rho}{\partial t} + \vec{\nabla} \cdot \vec{J} = 0$  ( )

(b)  $\frac{\partial \rho}{\partial t} - \vec{\nabla} \cdot \vec{J} = 0$  ( )

(c)  $\frac{\partial \rho}{\partial t} + \vec{\nabla} \cdot \vec{P} = 0$  ( )

(d)  $\frac{\partial \rho}{\partial t} - \vec{\nabla} \cdot \vec{P} = 0$  ( )

7. Let  $E_3$  be energy of the third energy level of a free particle in one-dimensional infinite potential well. The relation between first energy level  $E_1$  and third energy level  $E_3$  is

(a)  $E_3 = 3E_1$  ( )

(b)  $E_3 = E_1$  ( )

(c)  $E_1 = 9E_3$  ( )

(d)  $E_3 = 9E_1$  ( )

8. For a free particle in step potential, let  $R$  and  $T$  be reflectance and transmittance, then

(a)  $R + T = 1$  ( )

(b)  $R = T$  ( )

(c)  $R - T = 1$  ( )

(d)  $R \cdot T = 1$  ( )

9. The orbital magnetic moment of an electron is given (where  $L$  is angular momentum and  $m$  is mass of electron) by

(a)  $\mu_L = \frac{eL}{2m}$  ( )

(b)  $\mu_L = \frac{ne\hbar}{2m}$  ( )

(c)  $\mu_L = \frac{neL}{2m}$  ( )

(d) Both (a) and (b) ( )

10. For electron, the number of possible spin states for  $Z$  component is

(a) 1 ( )

(b) 2 ( )

(c) 3 ( )

(d) 4 ( )

SECTION—II

( Marks : 15 )

Give short answers to the following questions : 3×5=15

1. Show that a material particle cannot be equivalent to a single wave starting from de Broglie relation.

2. Show that  $[x, p_x^n] = n\hbar p_x^{n-1}$ , where  $x$  is position operator,  $p_x$  is  $x$  component of momentum operator.

Give short answers to the following questions:

1. Show that a material particle cannot be equivalent to a single wave starting from de Broglie relation.

9. The orbital magnetic moment of electron is given by  $\mu_L = \frac{e\hbar}{2m} \sqrt{L(L+1)}$  where  $L$  is the maximum value of  $l$  (orbital angular momentum) by

(a)  $\mu_L = \frac{e\hbar}{2m}$

(b)  $\mu_L = \frac{e\hbar}{m}$

(c)  $\mu_L = \frac{ne\hbar}{2m}$

- (d) Both (a) and (b)

10. For electron, the number of possible spin states for  $Z$  component is

(a) 1

(b) 2

(c) 3

(d) 4

3. What do you mean by eigenvalue and eigenfunction? Is the function  $e^{ax}$  an eigenfunction with respect to the operator  $\frac{d}{dx}$ ? If so, what are the eigenvalue and eigenfunction?

PHYSICS

SIXTH PAPER

( Quantum Mechanics - II )

( PART : A - OBJECTIVE )

( Marks : 25 )

The figures in the margin indicate full marks for the questions

SECTION - I

( Marks : 10 )

Put a Tick (✓) mark against the correct answer in the brackets provided :

1. Of the following particles moving with the same velocity, the one which has the largest wavelength is

(a) an electron

(b) a proton

(c) a neutron

(d) an  $\alpha$ -particle

4. The solution for a free particle in an infinitely high potential is given by  $\psi = A \sin kx + B \cos kx$ . If both the constants  $A$  and  $B$  are zero each, what physical meaning does it imply?

- (a) a wave packet
- (b) a light wave
- (c) Cannot behave as wave

3. Eigenvalues of Hermitian operators

- (a) are real only
- (b) are imaginary only
- (c) can be real or imaginary
- (d) are always complex

4. Let  $\vec{u} = (1, 2, -4)$  and  $\vec{v} = (2, -1, 3)$  be any two vectors in  $R^3$  space. Their inner product is equal to

- (a) 10
- (b) 7
- (c) -5
- (d) 15

5. Show that electron spin magnetic moment is equal to Bohr magnetron.

\*\*\*